

Technical Data Sheet: hTERT-immortalized White Preadipocytes

| | |
|------------------------------|---|
| ATCC® Number | CRL-4063™ |
| Organism | <i>Homo sapiens</i> |
| Tissue/Disease Source | Subcutaneous abdominal adipose tissue |
| Product Description | hTERT immortalized white preadipocytes were isolated from subcutaneous abdominal adipose tissue from a donor with von Hippel-Lindau syndrome. |
| Application | This hTERT-immortalized primary cell has applications as an in vitro cell model for toxicity studies and the study of obesity and related diseases. |




Figure 1: Cell morphology of Immortalized White Preadipocytes. Cells were maintained in ATCC recommended culture conditions. High and low confluence images of plated adherent white preadipocytes were taken using Nikon microscope at 10x. Scale bar represents 100 microns.

Preadipocyte Marker Analysis




Figure 2: Marker expression by flow cytometry. Cells were stained for CD90 and CD166 which are common markers for cells of mesenchymal stem cell origin, and CD34, a marker for hematopoietic stem cells. As expected, white preadipocytes were positive for CD90, CD166, but negative for CD34.

Differentiation of White Preadipocytes into Mature Adipocytes




Figure 3: Differentiation of hTERT White Preadipocytes from day 0 through day 25. Images show the morphology of differentiating cells at different timepoints.

Protocol:

- Culture cells to reach around 70% confluent, seed cells into 6 well plates at a seeding density of 100k cells/cm².
- Observe cells under a phase contrast microscope daily until cells are 100% confluent. Wait 48 additional hours to begin differentiation/induction.
- Replace the cell culture media with Induction Media 1. Incubate the cells for 7 days.
- Switch to Induction Media 2. Grow the cells for additional 14-18 days until Oil droplets can be seen clearly in the mature adipocytes.

Media Formulations:

Induction Media 1:

- Advanced DMEM
- 2% Sigma FBS
- 1% Pen/Strep
- 0.5 uM Human Insulin
- 0.1uM Dexamethasone
- 0.5 mM IBMX
- 33 uM Biotin
- 2 nM T3
- 30 uM Indomethacin
- 17 uM Pantothenate
- 2 uM Rosiglitazone

Induction Media 2:

- Advanced DMEM
- 2% Sigma FBS
- 1% Pen/Strep
- 0.5 uM Human Insulin
- 0.1uM Dexamethasone
- 0.5 mM IBMX
- 33 uM Biotin
- 2 nM T3
- 30 uM Indomethacin
- 17 uM Pantothenate